

Cholinesterase Testing Protocols for Healthcare Providers

Whom to Test?

Cholinesterase-inhibiting Pesticides

Test if working with Class I and Class II organophosphates (OP) or OP and N-methyl-carbamates for greater than a total of 30 hours in 30 consecutive days.

N-methyl-carbamates

If only working with N-methyl-carbamates, cholinesterase testing is not likely to be beneficial.

Baseline

Baseline Determination

Obtain baseline measures prior to working with cholinesterase-inhibiting pesticides. When obtaining the baseline, ensure that at least the previous 30 days were free of OP exposures.

Ensure that cholinesterase-inhibiting pesticides had not been handled in the immediate 30 days prior to testing.*

Establish baselines annually.

Working Baselines

Working baselines (baselines that are established when a 30-day period free of OP exposure is not possible) are likely to increase false negatives. Perform a second baseline after halting exposure (the longest practicable exposure-free period available is recommended, with a one-week exposure-free period at a minimum). If values differ by more than 10%, obtain a third baseline. The highest value should be used as the baseline.

2nd Baseline

A second baseline is recommended for improved precision but not essential. If a 2nd baseline is obtained, average the two values. When obtaining the 2nd baseline, wait to test until at least 3 days after the baseline, but within 30 days and ensure no pesticide exposures during this time period.

Testing

Test Types

Measure both acetylcholinesterase (red blood cell cholinesterase-AChE) and butyryl cholinesterase (plasma cholinesterase-PChE). AChE and PChE tests recommended; PChE if only performing 1 test.

Laboratory Services

Use the same laboratory and the same methodology for all testing so that results may be accurately compared.

Post Exposure Testing

Conduct post exposure test each time worker exceeds or reaches 30 hours of exposure within any 30-day period after the baseline or last post exposure test.

Medical Removal

Remove from handling cholinesterase-inhibiting pesticides with 30% or more reduction in cholinesterase activity (depression) of RBC or 40% or more reduction of plasma cholinesterase activity (depression).

Level to Return to Handling

Return to Handling

Return to handling when test result is greater than or equal to 80% of baseline.

Retest for Return to Work

Days to repeat test is determined by degree of reduction in cholinesterase activity.

For RBC AChE: $(\% \text{ depression} - 20) / 0.83 = \#$ of days to repeat test

For Plasma PChE: $(\% \text{ depression} - 20) / 1.2 = \#$ of days to repeat test

Review of Handling Practices

Review pesticide handling practices when test results are less than 80% of baseline.

* Handling of pesticides refers to tasks such as mixing, loading, transferring or applying pesticides; handling open containers of pesticides; acting as a flagger; cleaning, handling, adjusting or repairing pesticide equipment; or assisting with the application of pesticides.

Acknowledgements

These protocols were developed by *National Farm Medicine Center, AgriSafe Network and Migrant Clinicians Network*. **Matthew Keifer, MD, MPH** and **Carolyn Sheridan, RN, BSN** served as the lead authors with support from **Amy K. Liebman, MPA, MA**. The protocols are based on a review seven cholinesterase protocols and endorsed by MCN's Environmental and Occupational Health Advisory Committee and AgriSafe's Quality Assurance Committee.

REVIEWERS: Denise Andress, RN, BSN, Southwest Area Health Ed Center ▪ Geoffrey Calvert, MD, MPH, FACP, National Institute for Occupational Safety and Health ▪ Stephanie Chalupka, EdD, RN, PHCNS-BC, FAAOHN, Worcester State University ▪ Elizabeth Freeman Lambar, North Carolina Farmworker Health Program ▪ John Furman, PhD, MSN, CIC, COHN-S, Washington Department of Health, Susan Guin, MSN, University of Alabama ▪ Charlotte Halverson, BSN, COHN-S, National Education Center for Agricultural Safety ▪ Wilton C. Kennedy, DHSc, PA-C, MMSC, Jefferson College of Health Sciences ▪ Candace Kugel, CRNP, CNM, Migrant Clinicians Network ▪ James Roberts, MD, MPH, Medical University of South Carolina ▪ Lisa Schiller, PhD, APNP, FNP-BC, University of Wisconsin - Eau Claire ▪ Daniel L. Sudakin, MD, MPH, FACMT, FACOEM, Oregon State University ▪ Ed Zuroweste, MD, Migrant Clinicians Network

Sources

AgriSafe Network. February, 2007. *Standing Orders for Agricultural Occupational Screening Exam*. Spencer, IA: AgriSafe Network. <http://www.agrisafe.org/files/Standing%20Orders%202-23-07.pdf>

Brown A, Miller M and Keifer M. June, 2006. *Cholinesterase Monitoring – A Guide for the Health Professional*. College Park, MD: University of Maryland Cooperative Extension. Pesticide Information Leaflet No. 30.

Furman J. *Cholinesterase Monitoring for Agricultural Pesticide Handlers: Guidelines for Health Care Providers in Washington State*. Olympia, Washington: Washington State Department of Labor and Industries, Division of Occupational Safety and Health; January, 2010. <http://www.lni.wa.gov/Safety/Topics/AtoZ/Cholinesterase/files/ProvidersGuidelines1.pdf>

Ngai W, Ames R, Wisniewski J and Fan A. 2002. *Guidelines for Physicians who Supervise Workers Exposed to Cholinesterase-Inhibiting Pesticides*. Oakland, CA: California Environmental Protection Agency, Office of Environmental Health Hazard Assessment. 4th Edition. <http://oehha.ca.gov/pesticides/pdf/docguide2002.pdf>

Partners in Agricultural Health. 2003. *Cholinesterase Testing and Screening*. Madison, WI: Wisconsin Office of Rural Health. <http://worh.org/files/AgHealth/choltest.pdf>

Cholinesterase Health Monitoring Program for USDA Employees. (2004). *Cholinesterase Testing Program*. Washington, DC: U.S. Department of Agriculture: Animal, Plant and Health Inspection Service. http://www.aphis.usda.gov/emergency_response/downloads/health/Appendix%206%20B%20cholinesterase%20teating%20program.pdf

DISCLAIMER: This material was produced, in part, under Assistance Agreement No. X883487601 awarded by the U.S. Environmental Protection Agency. It has not been formally reviewed by EPA. EPA does not endorse any products or commercial services mentioned in this product. This material was produced, in part, under grant SH-22284-11-60-F-19 from the Occupational Safety and Health Administration, U.S. Department of Labor. It does not necessarily reflect the views or policies of the U.S. Department of Labor, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government. The project described was supported by Grant Number U30CS09742 from Health Resources and Services Administration. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Health Resources and Services Administration.

